Document - CODE DX Fountain

DX H66.90 DX-1 DX-2 CPT 99309 CPT-1 CPT-2 Chief Complaint (CC) HPI Allergies Levofloxacin, phenytoin, banana Current Medications ICD-10 Diagnosis H66.90 Otitis media, unspecified, unspecified ear Antibiotics Neomycin-polymyxin-HC solution; 3.5-10,000-1 mg/mL-unit/mL-%; amt: 4 gits tid; otic (ear) Special Instructions: administer neomycin-polymyxin-HC 4 gits on his left ear) q8 hours for 7 days. Otitis Media, unspecified H66.90 Every 8 Hours. Begin: 03/23/2025 - Open Ended Daily shower Oral care Change position every two hours Other recommendations Treatment should focus on symptom management and addressing potential underlying causes. If bacterial infection is suspected, a course of oral antibiotics (such as amoxicillin) may be appropriate, while analgesics like acetaminophen or NSAIDs can help alleviate pain. Ear drops may be considered if there is significant discomfort, but caution is needed to avoid exacerbating any underlying conditions. Encourage the patient to stay hydrated, practice ear hygiene, and avoid irritants like smoking. If symptoms persist or worsen, referral to an otolaryngologist may be necessary to rule out complications such as chronic otitis media or hearing loss. CPT code (s) Confirmed YES \$ 24 Count 0	Document - CODE DX Fountain	
DX-1 DX-2 CPT OPT 99309 CPT-1 CPT-2 Chief Complaint (CC) HPI Allergies Levofloxacin, phenytoin, banana Current Medications ICD-10 Diagnosis H66.90 Otitis media, unspecified, unspecified ear Antibiotics Neomycin-polymyxin-HC solution; 3.5-10,000-1 mg/mL-unit/mL-%; amt: 4 gtts tid; otic (ear) Special Instructions: administer neomycin-polymyxin-HC (4 gtts on his left ear) q8 hours for 7 days. Otitis Media, unspecified H66.90 Every 8 Hours. Begin: 03/23/2025 - Open Ended □ Ending: Open Ended □ Daily shower Oral care Change position every two hours Other recommendations Treatment should focus on symptom management and addressing potential underlying causes. If bacterial infection is suspected, a course of oral antibiotics (such as amoxicillin) may be appropriate, while analgesics like acetaminophen or NSAIDs can help alleviate pain. Ear drops may be considered if there is significant discomfort, but caution is needed to avoid exacerbating any underlying conditions. Encourage the patient to stay hydrated, practice ear hygiene, and avoid irritants like smoking. If symptoms persist or worsen, referral to an otolaryngologist may be necessary to rule out complications such as chronic otitis media or hearing loss. CPT code (s) Confirmed YES	NOTES	
DX-2 CPT 99309 CPT-1 CPT-2 Chief Complaint (CC) HPI Allergies Levofloxacin, phenytoin, banana Current Medications ICD-10 Diagnosis H66.90 Otitis media, unspecified, unspecified ear Antibiotics Neomycin-polymyxin-HC solution; 3.5-10,000-1 mg/mL-unit/mL-%; amt: 4 gits tid; otic (ear) Special Instructions: administer neomycin-polymyxin-HC (4 gits on his left ear) q8 hours for 7 days. Otitis Media, unspecified H66.90 Every 8 Hours. Begin: 03/23/2025 - Open Ended □ Ending: Open Ended □ Daily shower Oral care Change position every two hours Other recommendations Treatment should focus on symptom management and addressing potential underlying causes. If bacterial infection is suspected, a course of oral antibiotics (such as amoxicillin) may be appropriate, while analgesics like acetaminophen or NSAIDs can help alleviate pain. Ear drops may be considered if there is significant discomfort, but caution is needed to avoid exacerbating any underlying conditions. Encourage the patient to stay hydrated, practice ear hygiene, and avoid irritants like smoking. If symptoms persist or worsen, referral to an otolaryngologist may be necessary to rule out complications such as chronic otitis media or hearing loss. CPT code (s) Confirmed YES	DX	H66.90
CPT-1 CPT-2 Chief Complaint (CC) HPI Allergies Levofloxacin, phenytoin, banana Current Medications ICD-10 Diagnosis H66.90 Otitis media, unspecified, unspecified ear Antibiotics Neomycin-polymyxin-HC solution; 3.5-10,000-1 mg/mL-unit/mL-%; amt: 4 gtts tid; otic (ear) Special Instructions: administer neomycin-polymyxin-HC (4 gtts on his left ear) q8 hours for 7 days.Otitis Media, unspecified H66.90 Every 8 Hours. Begin: 03/23/2025 - Open Ended Daily shower Oral care Change position every two hours Other recommendations Treatment should focus on symptom management and addressing potential underlying causes. If bacterial infection is suspected, a course of oral antibiotics (such as amoxicillin) may be appropriate, while analgesics like acetaminophen or NSAIDs can help alleviate pain. Ear drops may be considered if there is significant discomfort, but caution is needed to avoid exacerbating any underlying conditions. Encourage the patient to stay hydrated, practice ear hygiene, and avoid irritants like smoking. If symptoms persist or worsen, referral to an otolaryngologist may be necessary to rule out complications such as chronic otitis media or hearing loss. CPT code (s) Confirmed YES	DX-1	
CPT-2 Chief Complaint (CC) HPI Allergies Levofloxacin, phenytoin, banana Current Medications ICD-10 Diagnosis H66.90 Otitis media, unspecified, unspecified ear Antibiotics Neomycin-polymyxin-HC solution; 3.5-10,000-1 mg/mL-unit/mL-%; amt: 4 gtts tid; otic (ear) Special Instructions: administer neomycin-polymyxin-HC (4 gtts on his left ear) q8 hours for 7 days.Otitis Media, unspecified H66.90 Every 8 Hours. Begin: 03/23/2025 - Open Ended □ Ending: Open Ended □ Daily shower Oral care Change position every two hours Other recommendations Treatment should focus on symptom management and addressing potential underlying causes. If bacterial infection is suspected, a course of oral antibiotics (such as amoxicillin) may be appropriate, while analgesics like acetaminophen or NSAIDs can help alleviate pain. Ear drops may be considered if there is significant discomfort, but caution is needed to avoid exacerbating any underlying conditions. Encourage the patient to stay hydrated, practice ear hygiene, and avoid irritants like smoking. If symptoms persist or worsen, referral to an otolaryngologist may be necessary to rule out complications such as chronic otitis media or hearing loss. CPT code (s) Confirmed YES \$ 24	DX-2	
CPT-2 Chief Complaint (CC) HPI Allergies Levofloxacin, phenytoin, banana Current Medications ICD-10 Diagnosis H66.90 Otitis media, unspecified, unspecified ear Antibiotics Neomycin-polymyxin-HC solution; 3.5-10,000-1 mg/mL-unit/mL-%; amt: 4 gtts tid; otic (ear) Special Instructions: administer neomycin-polymyxin-HC (4 gtts on his left ear) q8 hours for 7 days. Otitis Media, unspecified H66.90 Every 8 Hours. Begin: 03/23/2025 - Open Ended□ Ending: Open Ended Daily shower Oral care Change position every two hours Other recommendations Treatment should focus on symptom management and addressing potential underlying causes. If bacterial infection is suspected, a course of oral antibiotics (such as amoxicillin) may be appropriate, while analgesics like acetaminophen or NSAIDs can help alleviate pain. Ear drops may be considered if there is significant discomfort, but caution is needed to avoid exacerbating any underlying conditions. Encourage the patient ostay hydrated, practice ear hydgiene, and avoid irritants like smoking. If symptoms persist or worsen, referral to an otolaryngologist may be necessary to rule out complications such as chronic otitis media or hearing loss. CPT code (s) Confirmed YES	CPT	99309
Chief Complaint (CC) HPI Allergies Levofloxacin, phenytoin, banana Current Medications ICD-10 Diagnosis H66.90 Otitis media, unspecified, unspecified ear Antibiotics Neomycin-polymyxin-HC solution; 3.5-10,000-1 mg/mL-unit/mL-%; amt: 4 gtts tid; otic (ear) Special Instructions: administer neomycin-polymyxin-HC (4 gtts on his left ear) q8 hours for 7 days. Otitis Media, unspecified H66.90 Every 8 Hours. Begin: 03/23/2025 - Open Ended □ Daily shower Oral care Change position every two hours Other recommendations Treatment should focus on symptom management and addressing potential underlying causes. If bacterial infection is suspected, a course of oral antibiotics (such as amoxicillin) may be appropriate, while analgesics like acetaminophen or NSAIDs can help alleviate pain. Ear drops may be considered if there is significant discomfort, but caution is needed to avoid exacerbating any underlying conditions. Encourage the patient to stay hydrated, practice ear hygiene, and avoid irritants like smoking. If symptoms persist or worsen, referral to an otolaryngologist may be necessary to rule out complications such as chronic otitis media or hearing loss. CPT code (s) Confirmed YES \$ 24	CPT-1	
Allergies Levofloxacin, phenytoin, banana Current Medications ICD-10 Diagnosis H66.90 Otitis media, unspecified, unspecified ear Antibiotics Neomycin-polymyxin-HC solution; 3.5-10,000-1 mg/mL-unit/mL-%; amt: 4 gtts tid; otic (ear) Special Instructions: administer neomycin-polymyxin-HC (4 gts tid) on his left ear) q8 hours for 7 days. Otitis Media, unspecified H66.90 Every 8 Hours. Begin: 03/23/2025 - Open Ended □ Ending: Open Ended □ Daily shower Oral care Change position every two hours Other recommendations Treatment should focus on symptom management and addressing potential underlying causes. If bacterial infection is suspected, a course of oral antibiotics (such as amoxicillin) may be appropriate, while analgesics like acetaminophen or NSAIDs can help alleviate pain. Ear drops may be considered if there is significant discomfort, but caution is needed to avoid exacerbating any underlying conditions. Encourage the patient to stay hydrated, practice ear hygiene, and avoid irritants like smoking. If symptoms persist or worsen, referral to an otolaryngologist may be necessary to rule out complications such as chronic otitis media or hearing loss. CPT code (s) Confirmed YES \$ 24	CPT-2	
Current Medications ICD-10 Diagnosis H66.90 Otitis media, unspecified, unspecified ear Antibiotics Neomycin-polymyxin-HC solution; 3.5-10,000-1 mg/mL-unit/mL-%; amt: 4 gtts tid; otic (ear) Special Instructions: administer neomycin-polymyxin-HC (4 gtts on his left ear) q8 hours for 7 days. Otitis Media, unspecified H66.90 Every 8 Hours. Begin: 03/23/2025 - Open Ended Daily shower Oral care Change position every two hours Other recommendations Treatment should focus on symptom management and addressing potential underlying causes. If bacterial infection is suspected, a course of oral antibiotics (such as amoxicillin) may be appropriate, while analgesics like acetaminophen or NSAIDs can help alleviate pain. Ear drops may be considered if there is significant discomfort, but caution is needed to avoid exacerbating any underlying conditions. Encourage the patient to stay hydrated, practice ear hygiene, and avoid irritants like smoking. If symptoms persist or worsen, referral to an otolaryngologist may be necessary to rule out complications such as chronic otitis media or hearing loss. CPT code (s) Confirmed YES \$ 24	Chief Complaint (CC)	
Current Medications ICD-10 Diagnosis H66.90 Otitis media, unspecified, unspecified ear Antibiotics Neomycin-polymyxin-HC solution; 3.5-10,000-1 mg/mL-unit/mL-%; amt: 4 gtts tid; otic (ear) Special Instructions: administer neomycin-polymyxin-HC (4 gtts on his left ear) q8 hours for 7 days.Otitis Media, unspecified H66.90 Every 8 Hours. Begin: 03/23/2025 - Open Ended □ Ending: Open Ended □ Daily shower Oral care Change position every two hours Other recommendations Treatment should focus on symptom management and addressing potential underlying causes. If bacterial infection is suspected, a course of oral antibiotics (such as amoxicillin) may be appropriate, while analgesics like acetaminophen or NSAIDs can help alleviate pain. Ear drops may be considered if there is significant discomfort, but caution is needed to avoid exacerbating any underlying conditions. Encourage the patient to stay hydrated, practice ear hygiene, and avoid irritants like smoking. If symptoms persist or worsen, referral to an otolaryngologist may be necessary to rule out complications such as chronic otitis media or hearing loss. CPT code (s) Confirmed YES \$ 24	HPI	
ICD-10 Diagnosis H66.90 Otitis media, unspecified, unspecified ear Antibiotics Neomycin-polymyxin-HC solution; 3.5-10,000-1 mg/mL-unit/mL-%; amt: 4 gtts tid; otic (ear) Special Instructions: administer neomycin-polymyxin-HC (4 gtts on his left ear) q8 hours for 7 days.Otitis Media, unspecified H66.90 Every 8 Hours. Begin: 03/23/2025 - Open Ended □ Ending: Open Ended □ Daily shower Oral care Change position every two hours Treatment should focus on symptom management and addressing potential underlying causes. If bacterial infection is suspected, a course of oral antibiotics (such as amoxicillin) may be appropriate, while analgesics like acetaminophen or NSAIDs can help alleviate pain. Ear drops may be considered if there is significant discomfort, but caution is needed to avoid exacerbating any underlying conditions. Encourage the patient to stay hydrated, practice ear hygiene, and avoid irritants like smoking. If symptoms persist or worsen, referral to an otolaryngologist may be necessary to rule out complications such as chronic otitis media or hearing loss. CPT code (s) Confirmed YES \$ 24	Allergies	Levofloxacin, phenytoin, banana
Antibiotics Neomycin-polymyxin-HC solution; 3.5-10,000-1 mg/mL-unit/mL-%; amt: 4 gtts tid; otic (ear) Special Instructions: administer neomycin-polymyxin-HC (4 gtts on his left ear) q8 hours for 7 days.Otitis Media, unspecified H66.90 Every 8 Hours. Begin: 03/23/2025 - Open Ended ☐ Ending: Open Ended ☐ Daily shower Oral care Change position every two hours Other recommendations	Current Medications	
unit/mL-%; amt: 4 gtts tid; otic (ear) Special Instructions: administer neomycin-polymyxin-HC (4 gtts on his left ear) q8 hours for 7 days.Otitis Media, unspecified H66.90 Every 8 Hours. Begin: 03/23/2025 - Open Ended□ Ending: Open Ended Daily shower Oral care Change position every two hours Other recommendations Treatment should focus on symptom management and addressing potential underlying causes. If bacterial infection is suspected, a course of oral antibiotics (such as amoxicillin) may be appropriate, while analgesics like acetaminophen or NSAIDs can help alleviate pain. Ear drops may be considered if there is significant discomfort, but caution is needed to avoid exacerbating any underlying conditions. Encourage the patient to stay hydrated, practice ear hygiene, and avoid irritants like smoking. If symptoms persist or worsen, referral to an otolaryngologist may be necessary to rule out complications such as chronic otitis media or hearing loss. CPT code (s) Confirmed YES \$ 24	ICD-10 Diagnosis	H66.90 Otitis media, unspecified, unspecified ear
addressing potential underlying causes. If bacterial infection is suspected, a course of oral antibiotics (such as amoxicillin) may be appropriate, while analgesics like acetaminophen or NSAIDs can help alleviate pain. Ear drops may be considered if there is significant discomfort, but caution is needed to avoid exacerbating any underlying conditions. Encourage the patient to stay hydrated, practice ear hygiene, and avoid irritants like smoking. If symptoms persist or worsen, referral to an otolaryngologist may be necessary to rule out complications such as chronic otitis media or hearing loss. CPT code (s) YES \$ 24	Antibiotics	unit/mL-%; amt: 4 gtts tid; otic (ear) Special Instructions: administer neomycin-polymyxin-HC (4 gtts on his left ear) q8 hours for 7 days.Otitis Media, unspecified H66.90 Every 8 Hours. Begin: 03/23/2025 - Open Ended□ Ending: Open Ended Daily shower Oral care
Confirmed YES \$ 24	Other recommendations	addressing potential underlying causes. If bacterial infection is suspected, a course of oral antibiotics (such as amoxicillin) may be appropriate, while analgesics like acetaminophen or NSAIDs can help alleviate pain. Ear drops may be considered if there is significant discomfort, but caution is needed to avoid exacerbating any underlying conditions. Encourage the patient to stay hydrated, practice ear hygiene, and avoid irritants like smoking. If symptoms persist or worsen, referral to an otolaryngologist may be necessary to rule out complications
\$ 24	CPT code (s)	
	Confirmed	YES
Count 0	\$	24
	Count	0