Document - CODE DX Fountain

DOB	7 22 4050
SEX	7-22-1959
ENSURANCE	
NOTES	
Photos or/and Videos	M96 O
DX	M86.9
DX-1	
DX-2	00000
CPT	99308
CPT-1	
CPT-2	
Chief Complaint (CC)	Stable for removal of hardware. Monitor for Cephalexin antibiotic. Care team continue to express concerns regarding the patient's frailty and chronic health deterioration.
HPI	The patient is 65-year-old male, with PMH of Osteomyelitis, unspecified, Charcot's joint, right ankle and foot, Valgus deformity, not elsewhere classified, unspecified ankle, Peripheral vascular disease, unspecified, Lymphedema, not elsewhere classified, Essential (primary) hypertension, Other psychoactive substance abuse, uncomplicated. The patient is lying down; remains in stable condition from an ID standpoint. No acute distress reported by nursing staff. He denies chest pain, sob or palpitations. Patient is AAOX1. Unable to actively participate in their care due to underlying conditions. No family at bedside. Medication review in chart. Tolerating Cephalexin antibiotic medications well. V/S reviewed and are WNL. Stable for removal of hardware. Care team continue to express concerns regarding the patient's frailty and chronic health deterioration.
Allergies	No known allergies
	3 3 3
Current Medications	Cephalexin capsule; 500 mg; amt: 500 mg; oral Special Instructions: Dx: Prophylaxis for Removal of Hardware Every 8 Hours. Begin: 04/05/2025 - 04/14/2025 Ending: 04/14/2025 Gabapentin capsule; 300 mg; amt: 300 mg; oral Special Instructions: Dx: neuropathic pain Three Times A Day. Begin: 04/05/2025 - Open ended.
Current Medications ICD-10 Diagnosis	Cephalexin capsule; 500 mg; amt: 500 mg; oral Special Instructions: Dx: Prophylaxis for Removal of Hardware Every 8 Hours. Begin: 04/05/2025 - 04/14/2025 Ending: 04/14/2025 Gabapentin capsule; 300 mg; amt: 300 mg; oral Special Instructions: Dx: neuropathic pain Three Times A Day. Begin:
	Cephalexin capsule; 500 mg; amt: 500 mg; oral Special Instructions: Dx: Prophylaxis for Removal of Hardware Every 8 Hours. Begin: 04/05/2025 - 04/14/2025 Ending: 04/14/2025 Gabapentin capsule; 300 mg; amt: 300 mg; oral Special Instructions: Dx: neuropathic pain Three Times A Day. Begin: 04/05/2025 - Open ended.
ICD-10 Diagnosis	Cephalexin capsule; 500 mg; amt: 500 mg; oral Special Instructions: Dx: Prophylaxis for Removal of Hardware Every 8 Hours. Begin: 04/05/2025 - 04/14/2025 Ending: 04/14/2025 Gabapentin capsule; 300 mg; amt: 300 mg; oral Special Instructions: Dx: neuropathic pain Three Times A Day. Begin: 04/05/2025 - Open ended. M86.9 Osteomyelitis, unspecified Cephalexin capsule; 500 mg; amt: 500 mg; oral Special Instructions: Dx: Prophylaxis for Removal of Hardware Every 8 Hours. Begin: 04/05/2025 - 04/14/2025 Ending: 04/14/2025 Gabapentin capsule; 300 mg; amt: 300 mg; oral Special Instructions: Dx: neuropathic pain Three Times A Day. Begin: 04/05/2025 - Open ended.
ICD-10 Diagnosis Antibiotics	Cephalexin capsule; 500 mg; amt: 500 mg; oral Special Instructions: Dx: Prophylaxis for Removal of Hardware Every 8 Hours. Begin: 04/05/2025 - 04/14/2025 Ending: 04/14/2025 Gabapentin capsule; 300 mg; amt: 300 mg; oral Special Instructions: Dx: neuropathic pain Three Times A Day. Begin: 04/05/2025 - Open ended. M86.9 Osteomyelitis, unspecified Cephalexin capsule; 500 mg; amt: 500 mg; oral Special Instructions: Dx: Prophylaxis for Removal of Hardware Every 8 Hours. Begin: 04/05/2025 - 04/14/2025 Ending: 04/14/2025 Gabapentin capsule; 300 mg; amt: 300 mg; oral Special Instructions: Dx: neuropathic pain Three Times A Day. Begin: 04/05/2025 - Open ended. Ending: Open ended It is recommended to ensure proper wound care and hygiene if there is an open site, with regular monitoring for signs of infection such as redness, swelling, warmth, or drainage. Offloading strategies Should be implemented if the infection involves weight-bearing bones, to reduce pressure and promote healing. Encourage adequate nutrition to support tissue repair, and consider physical therapy or mobility support as needed to prevent deconditioning. Ongoing assessment of pain, function, and any changes in the affected area is
ICD-10 Diagnosis Antibiotics Other recommendations	Cephalexin capsule; 500 mg; amt: 500 mg; oral Special Instructions: Dx: Prophylaxis for Removal of Hardware Every 8 Hours. Begin: 04/05/2025 - 04/14/2025 Ending: 04/14/2025 Gabapentin capsule; 300 mg; amt: 300 mg; oral Special Instructions: Dx: neuropathic pain Three Times A Day. Begin: 04/05/2025 - Open ended. M86.9 Osteomyelitis, unspecified Cephalexin capsule; 500 mg; amt: 500 mg; oral Special Instructions: Dx: Prophylaxis for Removal of Hardware Every 8 Hours. Begin: 04/05/2025 - 04/14/2025 Ending: 04/14/2025 Gabapentin capsule; 300 mg; amt: 300 mg; oral Special Instructions: Dx: neuropathic pain Three Times A Day. Begin: 04/05/2025 - Open ended. Ending: Open ended It is recommended to ensure proper wound care and hygiene if there is an open site, with regular monitoring for signs of infection such as redness, swelling, warmth, or drainage. Offloading strategies Should be implemented if the infection involves weight-bearing bones, to reduce pressure and promote healing. Encourage adequate nutrition to support tissue repair, and consider physical therapy or mobility support as needed to prevent deconditioning. Ongoing assessment of pain, function, and any changes in the affected area is important to guide care and prevent complications.
ICD-10 Diagnosis Antibiotics Other recommendations CPT code (s)	Cephalexin capsule; 500 mg; amt: 500 mg; oral Special Instructions: Dx: Prophylaxis for Removal of Hardware Every 8 Hours. Begin: 04/05/2025 - 04/14/2025 Ending: 04/14/2025 Gabapentin capsule; 300 mg; amt: 300 mg; oral Special Instructions: Dx: neuropathic pain Three Times A Day. Begin: 04/05/2025 - Open ended. M86.9 Osteomyelitis, unspecified Cephalexin capsule; 500 mg; amt: 500 mg; oral Special Instructions: Dx: Prophylaxis for Removal of Hardware Every 8 Hours. Begin: 04/05/2025 - 04/14/2025 Gabapentin capsule; 300 mg; amt: 300 mg; oral Special Instructions: Dx: neuropathic pain Three Times A Day. Begin: 04/05/2025 - Open ended. Ending: Open ended It is recommended to ensure proper wound care and hygiene if there is an open site, with regular monitoring for signs of infection such as redness, swelling, warmth, or drainage. Offloading strategies Should be implemented if the infection involves weight-bearing bones, to reduce pressure and promote healing. Encourage adequate nutrition to support tissue repair, and consider physical therapy or mobility support as needed to prevent deconditioning. Ongoing assessment of pain, function, and any changes in the affected area is important to guide care and prevent complications.

Count	1
COMPLETED	